

Critical Issues In Managing Pain With Opioid Therapy

How to Receive CME Credit

To earn free CME credit for watching this program, you must complete this test online.

Go to www.discoverychannelCME.com and follow the instructions to complete the post-test and evaluation online. You must be registered and signed-in in order to take the test online. Upon successful completion of the test and the evaluation, you will be asked to select your professional category and you will be able to print the appropriate credit certificate right from your computer.

CME Test Questions

1. Patients with a history of aberrant drug-related behaviors should never be considered for opioid therapy for the treatment of pain.
 - a. True
 - b. False

2. Data suggest that the incidence of addiction in the total pain patient population is _____%.
 - a. 1 to 2
 - b. 2 to 5
 - c. 10 to 20
 - d. 25 to 30
 - e. 50 to 60

3. According to SAMHSA, the number 1 source for pain medications for non-medical use among past year users is which of the following?
 - a. Free from a friend or family member
 - b. Internet
 - c. From one doctor
 - d. Bought from a friend or family member
 - e. Bought from a drug dealer

4. Which statement best describes prescription drug misuse?
 - a. State where patient without pain demonstrates behaviors consistent with addiction
 - b. State where patient with uncontrolled pain seeks medical care but may demonstrate behaviors of an addict
 - c. Tampering with a prescription drug to accelerate its effect
 - d. Taking a friend's or family member's medication to treat your pain
 - e. Taking your own medication not as prescribed

5. Goals of chronic opioid therapy include all of the following except:
 - a. Reduce pain intensity
 - b. Restore functional capacity
 - c. Deter misuse or abuse of opioids
 - d. Improve restful sleeping
 - e. Improve quality of life

6. An abuse-deterrent formulation may deter which type of abuse?
 - a. Crushing
 - b. Snorting
 - c. Chewing
 - d. Injecting
 - e. All of the above